

AD-A202 925

DTIC FILE COPY

AD _____

WORLD REFERENCE CENTER FOR ARBOVIRUSES

FINAL REPORT

Robert E. Shope, M.D.

January 1988

(For the period September 1, 1983 to August 31, 1986)

Supported by

U.S. Army Research and Development Command, Fort Detrick,
Frederick, Maryland 21701-5012

Grant No. DAMD17-83-G-9551

Yale University School of Medicine
New Haven, Connecticut 06510

DTIC
ELECTE
S JAN 26 1988
9 H

Approved for public release; distribution unlimited.

The views, opinions, and/or findings contained in this report are those of the author and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

89 1 26 049

SUMMARY:

The World Reference Center for Arboviruses maintains on a national and an international basis: a) serologic identification and biologic characterization of arboviruses using CF, HI, neutralization, IFA, and ELISA techniques and reagents prepared in large part in mice; b) molecular characterization of viruses using PAGE, electron microscopy, and polypeptide and RNA purification; c) diagnosis of disease using sera and other specimens submitted by the military and other organizations; d) diagnosis and epidemiological study of epidemics and epizootics using submitted specimens; e) preparation and distribution of reference reagents including antibody, viruses to specific organizations, and antigens under special circumstances; f) serological survey for arboviruses on a limited scale; and g) dissemination of information through WHO and the American Committee on Arthropod-borne Viruses. The above functions of the reference center were jointly supported by this grant and by grants from the U.S. NIH, the Australian government, and WHO.

During the 3 years of this grant, 197 viruses were identified, many of them new to science and/or new to a geographic region. The taxonomy of the rhabdoviruses, orbiviruses, and phleboviruses was revised. The distribution of antibody to arboviruses was determined throughout specific areas of the world by serosurvey. Techniques for rapid and early diagnosis of arbovirus diseases were refined and transferred as technology to users internationally. Two thousand, two hundred fifty-eight ampoules of arbovirus reagents were distributed to the U.S. Military and to other laboratories throughout the world.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

FOREWORD

In conducting the research described in this report, the investigator adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (DHEW Publication No. (NIH) 86-23, Revised 1985).

For the protection of human subjects the investigators have adhered to policies of applicable Federal Law 45CFR46.

BODY OF REPORT:

Introduction. The World Reference Center for Arboviruses was established at the Yale Arbovirus Research Unit in 1965 as an outgrowth of The Rockefeller Foundation program on arboviruses which was moved in 1965 to Yale University from New York City. The U.S. Army has supported this program since 1972, initially through joint Navy-Army funding, then through a separate contract, and during the past three years by this grant. The progress of the past three years is included in this report; it covers the work for the entire project which received support from WHO, NIH, and the Australian government, in addition to that of this grant.

Virus identification. A primary function of the reference center is to receive viruses from all parts of the world for identification. Attempts to identify about 200 isolates in mosquito cells from Indonesia determined that at least 64 of these were doublestranded segmented RNA viruses. Eleven of these had 9 segments, 8 had 10 segments in a pattern of 2-2-2-2-2, 11 had 10 segments similar to Matsu virus 2-4-3-1, one had 10 segments 3-3-3-1, 6 were EHD-like, 16 had 10 segments 3-6-1, 10 had 12 segments in a pattern 6-6, and one had 12 segments in a pattern 6-5-1. Several appeared to be mixtures with 3- or 5-segmented viruses as the second virus. Ten also reacted with the flavivirus grouping antibody, and 3 with alphavirus grouping antibody. Another Indonesian NAMRU isolate was a rhabdovirus, apparently new to science.

Viruses were identified from the United States and from many other nations as follows:

ALPHAVIRUSES. Chikungunya virus was identified from Culex mosquitoes from Central African Republic, and several strains of Highlands J virus were identified from Connecticut.

FLAVIVIRUSES. Dengue-4 was recognized from New Caledonia, several strains of Japanese encephalitis from Viet Nam, and a flavivirus from ticks of Connecticut (possibly a contaminant).

BUNYAVIRUSES. A new virus from ticks from India was in the Tete serogroup. Inkoo virus was identified for the first time from Sweden. A new Simbu group virus was recognized from Brazil and for the first time, a Bunyamwera serogroup virus was identified from Australia.

PHLEBOVIRUSES. New phleboviruses were characterized from Brazil and Colombia; Chagres was isolated for the first time from Colombian sandflies; and Toscana virus was identified from the cerebrospinal fluid of a tourist visiting in Portugal. Naples virus was identified, isolated by NAMRI-3 personnel in Egypt, and Sicilian and Naples-like viruses were identified from Swedish UN troops.

NAIROVIRUSES. The Xinjiang hemorrhagic fever virus from western China was identified as Crimean-Congo hemorrhagic fever virus. A strain of Avalon virus was found in Canada and a new nairovirus from a Brazilian bird was identified.

UUKUVIRUSES. Three new Uukuniemi serogroup viruses were characterized, Precarious Point, murre, and an as yet unnamed Alaskan isolate.

VESICULOVIRUSES. New VSV-related rhabdoviruses were recognized from Argentina and the United States. Also, a Tibrogargan-related rhabdovirus from Florida was characterized.

ORBIVIRUSES. In addition to the orbiviruses described above from Indonesia, several isolates of tick-borne (Kemerovo serogroup) viruses were identified from Canada, Australia, Finland, France, and California.

ARENAVIRUSES. Confirmation of the relationship of Ippy virus to Lassa virus was made. This finding was first reported from the Institute of Virology in South Africa.

UNCLASSIFIED VIRUSES. Several ungrouped and unclassified viruses were studied including Lake Clarendon virus from Australia, a virus from New York State voles, tick agents from western USA and from Tanzania, and a new mosquito isolate from Brazil.

Classification of arboviruses. The Quarantfil serogroup was characterized by electron microscopy as arenavirus-like. The serological relationships of a new member of the group were determined, and six proteins were found by PAGE analysis of infected cells. No serological relationship was found to other arenaviruses.

The members of the VSV serogroup were characterized by ELISA, neutralization, CF and immunofluorescence. The study included two new vesiculoviruses.

A comprehensive study of the orbiviruses was carried out including members of the bluetongue, EHD, Palyam, Corriparta, Kemerovo, and Colorado tick fever serogroups. The study embraced PAGE analyses and RNA-RNA blot hybridization. Extensive attempts to produce reassortants among orbiviruses showed that members of the serogroups reassorted, but that members of different serogroups did not reassort, even under extreme conditions. It was proposed that viruses which reassort are members of a single species, and that therefore viruses in a serogroup constituted a species. The RNA-RNA blot hybridization was demonstrated to be a sensitive and reliable identification method.

Diagnosis of disease. Sandfly fever was diagnosed in Swedish soldiers in Cyprus; EEE was confirmed in Whooping Cranes in Maryland; and equine, human, and avian encephalitis was identified in Connecticut.

Serologic and antigen surveys. Rift Valley fever antibody was shown in an indigenous population inhabiting the Senegal River Valley at the site near the river mouth where the Diama Dam has been constructed. Antibody to RVF was also shown in Zambia in farm residents who survived a hemorrhagic fever outbreak. The ELISA was used also to show widespread RVF antibody in sheep

and cattle from Mali. Antigen and antibody to a Hantaan-related virus was detected in voles from Connecticut. An antibody survey of Uganda revealed alpha-, flavi-, and bunyavirus reactivity.

Characterization of monoclonal antibodies. Sandfly Sicilian monoclonal antibodies supplied by scientists at Fort Detrick were characterized for their reactivity by immunofluorescence with a battery of phleboviruses.

Development of new techniques. The 4G2-4-15 dengue-2 monoclonal antibody was developed and used extensively as a universal flavivirus capture antibody in ELISA. Grouping monoclonal antibodies from CDC Fort Collins and from the University of Maryland were characterized and shown to be highly group reactive by HI and ELISA. The ELISA using SDS treated orbivirus antigen coat was developed for testing seabirds for Kemerovo group antibodies. An ELISA was elaborated for detecting Crimean-Congo hemorrhagic fever antibody in sheep, cattle, and humans. The technique was tested in Israel where CCHF antibody was found for the first time there in cattle. The enzyme immunoassay was adapted to use dengue virus-infected cultured mosquito cells as antigen and an arbovirus tissue culture neutralization test was developed using ELISA as an indicator. Mice immunized by intrasplenic inoculation with arboviruses developed ELISA antibody in 5 days. This method allows rapid reciprocal definitive identification of arboviruses.

Experimental pathogenesis. Hamsters were used as a model for testing viremia and immune response to sequential infections with phleboviruses. Immune enhancement was not shown, but broadening of immune response following sequential infections with different phleboviruses was evident.

Collection of low passage arbovirus strains. A large collection of low passage arbovirus strains was developed and maintained lyophilized. Priority was given to yellow fever, dengue, chikungunya, California encephalitis, Venezuelan encephalitis, St. Louis encephalitis, western encephalitis, eastern encephalitis, Japanese encephalitis, and other human disease arboviruses. Vesiculovirus strains of veterinary importance were also accessioned. The original (or as close to original as was available) material was passaged once in C6/36 mosquito cells or in Vero cells. The resulting stock was lyophilized in aliquots. These were stored and distributed to any and all persons requesting material for study.

Distribution of reagents. Reference antibodies and viruses were maintained as lyophilized stock for distribution to U.S. military users and to others on a world-wide basis. During the three years of the grant, 2,258 ampoules of arbovirus reagents were distributed to laboratories in more than 22 countries. This total consisted of 627 ampoules of virus stock, 1,055 ampoules of virus antigen, and 576 ampoules of antibody. The reagents distributed represented more than 200 different arboviruses.

Eight different cell lines were also distributed. These included both vertebrate and mosquito cells.

PUBLICATIONS

- Gligic, A., Tesh, R.B., Miscevic, Z., Travassos da Rosa, A.P.A. and Zivkovic, V. Jug Bogdanovac virus - a new member of the vesicular stomatitis virus serogroup (Rhabdoviridae: Vesiculovirus) isolated from phlebotomine sandflies in Yugoslavia. *Mikrobiologija* 20:97-105, 1983. (Published in 1984).
- Chastel, C., Main, A.J., Couatarmanc'h, A., LeLay, G., Guiguen, C., Knudson, D.L., Quillien, M.C. and Beaucournu, J.C. Isolation of Eyach virus from Ixodes ricinus and I. ventralloi ticks in France. *Arch. Virol.* 82:161-171, 1984.
- Knudson, D.L., Tesh, R.B., Main, A.J., St. George, T.D. and Digoutte, J.P. Characterization of the Palyam virus serogroup (Reoviridae: Orbivirus). *Intervirology* 22:41-49, 1984.
- Main, A.J. Eastern equine encephalomyelitis; Western equine encephalomyelitis; and Powassan virus *In* Geographical Distribution of Animal Viral Diseases, S. Odend'hal (ed.), Academic Press, 1984.
- Main, A.J., and Grans, W.J. Cache Valley virus from Aedes sollicitans in New Jersey, *J. Am. Mosq. Contr. Assoc.* 1984.
- Niklasson, B., Espmark, A., LeDuc, J.W., Gargan, T.P., Ennis, W.A., Tesh, R.B., and Main, A.J., Jr. Association of Sindbis virus with Ockelbo disease in Sweden. *Am. J. Trop. Med. & Hyg.*, 33:1212-1217, 1984.
- Scott, T.W. and Burrage, T.G. Rapid infection of salivary glands in Culiseta melanura with eastern equine encephalitis virus: an electron microscopic study. *Am. J. Trop. Med. & Hyg.* 33:961-964, 1984.
- Scott, T.W., Hildreth, S.W. and Beaty, B.J. The distribution and development of eastern equine encephalitis virus in its enzootic mosquito vector, Culiseta melanura. *Am. J. Trop. Med. & Hyg.* 34:300-310, 1984.
- Shope, R.E. Lyme disease: epidemiologic studies. *Yale J. Biol. Med.* 57:707, 1984.
- Shope, R.E. Currently available vaccines for Japanese encephalitis, and future research. *In* Dalrymple, J.M. (ed.) International Workshop on the Molecular Biology of Flaviviruses, p.25, U.S. Army Medical Research & Development Command, Fort Detrick, Frederick, MD, 1984.
- Shope, R.E. Japanese encephalitis. *In* Trevino, G.S. and Hyde, J.L. (eds.) Foreign Animal Diseases, 4th edition, pp. 216-222, U.S. Animal Health Association, Richmond, VA, 1984.
- St. George, T.D., Cybinski, D.H., Main, A.J., McKilligan, N., and Kemp, D.H. The isolation of a new arbovirus from Argas robertsi from a cattle egret colony in Australia. *Aus. J. Biol. Sci.* 37:85-89, 1984.
- Tesh, R.B. Transovarial transmission of arboviruses in their arthropod vectors, *In* Current Topics in Pathogen-Vector-Host Research, K.G. Harris (ed.) Praeger Scientific, New York, 2:57-76, 1984.

Tesh, R.B. and Modi, G.B. Studies on the biology of phleboviruses in sand flies (Diptera; Psychodidae). 1. Experimental infection of the vector. Am. J. Trop. Med. & Hyg. 33:1007-1016, 1984.

Tidwell, M.A., Forsythe, D.M., Parker, R.L. and Main, A.J. Eastern equine encephalomyelitis virus activity in South Carolina. J. Agr. Ent. 1:43-52, 1984.

Travassos da Rosa, A.P.A., Tesh, R.B., Pinheiro, F.P., Travassos da Rosa, J.F., Peralta, P.H. and Knudson, D.L. Characterization of the Changuinola virus serogroup (Reoviridae: Orbivirus). Intervirology 21:38-49, 1984.

Travassos da Rosa, A.P.A., Tesh, R.B., Travassos da Rosa, J.F. and Main A.J. Carajas and Maraba viruses, two new vesiculoviruses isolated from phlebotomine sandflies in Brazil. Am. J. Trop. Med. & Hyg., 33:999-1005, 1984.

Bodkin, D.K. and Knudson, D.L. Assessment of sequence relatedness of dsRNA genes by RNA-RNA blot hybridization. J. Virological Methods 10:45-52, 1985.

Bodkin, D.K. and Knudson, D.L. Sequence relatedness of Palyam virus genes to cognates of the Palyam serogroup viruses by RNA-RNA blot hybridization. Virology 143:55-62, 1985.

Chastel, C., Main, A.J., Guiguen, C., LeLay, G., Quillien, M.C., Monnat, J.Y., and Beaucournu, J.C. The isolation of Meaban virus, a new flavivirus from seabird tick Ornithodoros (Alectorobius) maritimus in France. Arch. Virol. 8:1-11, 1985.

Holford, T.R., Brown, S.E. and Knudson, D.L. Estimation of DNA fragment size and generation of DNA restriction endonuclease maps using linear models. J. Virol. Methods 10:117-126, 1985.

Gorman, B.M., Ghalib, H.W., Huismans, H., Joklik, W.K., Knudson, D.L., McPhee, D.A., Murphy, F.A., Ritchie, A., Roy, P., and Taylor, J. WHO/FAO working team report: Molecular virology, pp. 255-266, In Bluetongue and Related Orbiviruses, T.L. Barber and M.M. Jochim (eds.) Alan R. Liss, New York, 1985.

Knudson, D.L. and Shope, R.E. Overview of the orbiviruses. In Bluetongue and Related Orbiviruses (T.L. Barber and M.M. Jochim, eds.) pp. 255-256, Alan R. Liss, NY, 1985.

Rodhain, R., Metselaar, D., Ardoin, P., Hannoun, C., Shope, R.E. and Casals, J. Le virus Omo: Un nouveau virus du groupe Qalyub (Bunyaviridae, Nairovirus) isole de rongeurs in Ethiopie meridionale, Ann. Inst. Pasteur/Virol. 136E:243-247, 1985.

Shope, R.E. Arbovirus surveillance. In Surveillance of Viral Disease, B.D. Shoub (ed.) pp. 55-62, Poliomyelitis Research Foundation, Johannesburg, Republic of South Africa, 1985.

Shope, R.E. Viral spread between hosts, In Genetically Altered Viruses and the Environment, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, pp. 61-68, 1985.

St. George, T.D., Doherty, R.L., Carley, J.G., Filippich, C., Brescia, A., Casals, J., Kemp, D.H. and Brothers, N. The isolation of arboviruses including a new flavivirus and a new bunyavirus from Ixodes (Ceratiixodes) uriae (Ixodoidea: Ixodidae) collected at Macquarie Island, Australia, 1975-1979, Am. J. Trop. Med. & Hyg. 34:406-412, 1985.

Tesh, R.B. Undifferentiated fevers: dengue, phlebotomus fever, Rift Valley fever, West Nile fever and fevers caused by alphaviruses, pp. 1737-1740 In Cecil's Textbook of Medicine, 17th edition. J.B. Wyngaarden, L.H. Smith, and F. Plum (eds.) W.B. Saunders Co., Philadelphia, 1985.

Yen, Y.C., Kong, L.X., Lee, L., Zhang, Y.Q., Li, F., Cai, B.J., and Gao, S.Y. Characteristics of Crimean-Congo hemorrhagic fever virus (Xinjiang strain) in China. Am. J. Trop. Med. & Hyg. 34:1179-1182, 1985.

Bodkin, D.K. and Knudson, D.L. Genetic relatedness of Palyam serogroup viruses by RNA-RNA blot hybridization. J. Gen. Virol. 67:683-691, 1986.

Crans, W.J., McNelly, J., Schulze, T., and Main, A.J. Isolation of eastern equine encephalitis virus from Aedes sollicitans during an epizootic in southern New Jersey. J. Am. Mosq. Contr. Assoc., 2:68-72, 1986.

Kalunda, M., Mukwaya, L.G., Mukuye, A., Zale, M., Sekyalo, E., Wright, J. and Casals, J. Kasokero virus: A new human pathogen from bats (Rousettus aegyptiacus) in Uganda. Am. J. Trop. Med. & Hyg. 35:387-392, 1986.

Main, A.J. The bimodal seasonality of immature Ixodes dammini. J. Med Ent. (letter to the editor) 23:339, 1986.

Main, L.H. and Main, A.J. The potential health hazard of tires used in playgrounds. J. Nat. Assoc. Ed, Young Child. (letter to the editor) 41:159, 1986.

Olson, J.G., Ksiazek, T.G., Lee, V.H., Tan, R. and Shope, R.E. Isolation of Japanese encephalitis virus from Anopheles annularis and Anopheles vagus in Lombok, Indonesia. Trans. Roy. Soc. Trop. Med. Hyg. 79:845-847, 1986.

Rodhain, F., Leger, N., Madulo-LeBlond, Hannoun, C. and Tesh, R.B. Le virus Corfiou: un nouveau Phlebovirus isole de phlebotomes en Grece. Ann. Virol. 1986 in press.

Shope, R.E. Infectious disease research, In Viral and Mycoplasmal Infections of Laboratory Rodents, Chapter 20, pp. 679-687, P.N. Bhatt (ed.) Academic Press, NY, 1986.

St. George, T.D., Doherty, R.L., Carley, J.C., Filippich, C., Brescia, A., Casals, J., Kemp, D.H. and Brothers, N. The isolation of arboviruses including a new flavivirus and a new bunyavirus from Ixodes (Ceratiixodes) uriae (Ixodoidea: Ixodidae) collected at Macquarie Island, Southern Ocean 1975-1979. Am. J. Trop. Med. & Hyg. 34:406-412, 1985.

Tesh, R.B., Boshell, J., Young, D., Morales, A.A., Corredor, A.A., Modi, G., Ferro de Carrasquilla, C., de Rodriguez, C., and Gaitan, M.O. Biology of Arboledas virus, a new phlebotomus fever serogroup virus (Bunyaviridae: Phlebovirus) isolated from sand flies in Colombia. Am. J. Trop. Med. & Hyg. 35:1310-1316, 1986.

Tesh, R.B., Peleg, J., Samina, I., Margalit, J., Bodkin, D.K., Shope, R.E. and Knudson, D.L. Biological and antigenic characterization of Netivot virus, an unusual new Orbivirus recovered from mosquitoes in Israel. Am. J. Trop. Med. & Hyg. 35:418-428, 1986.

PROFESSIONAL PERSONNEL RECEIVING GRANT SUPPORT:

Robert E. Shope, M.D.
Susan Brown, Ph.D.
Andrew J. Main, Dr.P.H.
Dennis L. Knudson, D.Phil.

DISTRIBUTION LIST

12 copies	Director Walter Reed Army Institute of Research Walter Reed Army Medical Center ATTN: SGRD-UWZ-C Washington, DC 20307-5100
1 copy	Commander US Army Medical Research and Development Command ATTN: SGRD-RMI-S Fort Detrick, Frederick, Maryland 21701-5012
12 copies	Defense Technical Information Center (DTIC) ATTN: DTIC-DDAC Cameron Station Alexandria, VA 22304-6145
1 copy	Dean School of Medicine Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, MD 20814-4799
1 copy	Commandant Academy of Health Sciences, US Army ATTN: AHS-CDM Fort Sam Houston, TX 78234-6100

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188
Exp Date Jun 30 1986

1a REPORT SECURITY CLASSIFICATION Unclassified		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION Yale Univ. School of Medicine	6b OFFICE SYMBOL (If applicable)	7a NAME OF MONITORING ORGANIZATION	
6c ADDRESS (City, State, and ZIP Code) New Haven, Connecticut 06510		7b ADDRESS (City, State, and ZIP Code)	
8a NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command	8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-83-G-9551	
8c ADDRESS (City, State, and ZIP Code) Fort Detrick, Frederick, MD 21701-5012		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO 61102A	PROJECT NO 3M1 61102BS10
		TASK NO AC	WORK UNIT ACCESSION NO 073
11 TITLE (Include Security Classification) World Reference Center for Arboviruses (U)			
12 PERSONAL AUTHOR(S) Robert E. Shope, M.D.			
13a TYPE OF REPORT Final	13b TIME COVERED FROM 9/1/83 TO 8/31/86	14 DATE OF REPORT (Year, Month, Day) 1988 January	15 PAGE COUNT 12
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
06	13		Arbovirus, serological diagnosis, ELISA, complement-fixation
06	03		neutralization, serosurvey, Rift Valley fever, Crimean-Congo hemorrhagic fever, rapid diagnosis, flavivirus. (SDW)
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The World Reference Center for Arboviruses identified 197 viruses and revised the taxonomy of the Quarantil serogroup, rhabdoviruses, phleboviruses, and orbiviruses. Techniques for rapid and early diagnosis of arbovirus disease were refined and transferred as technology to users internationally. Virus reference reagents (2,258 ampoules) of over 200 viruses were distributed world-wide. <i>Keywords:</i>			
20 DISTRIBUTION STATEMENT <input checked="" type="checkbox"/> UNCLASSIFIED UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USER		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Mary Frances Bostian		22b TELEPHONE (Include Area Code) 301/663-7325	22c OFFICE SYMBOL SGRD-RMI-S